

# Help us figure out the best dose of aspirin!

We are recruiting patients that meet the following criteria for the ADAPTABLE Aspirin Study.

## INCLUSION CRITERIA ( $\geq 1$ ):

- Prior MI
- Prior PCI or CABG
- Prior angiogram showing significant CAD
- History of chronic ischemic heart disease, CAD, or ASCVD

## $\geq 1$ ENRICHMENT FACTOR:

- Age  $\geq 65$
- Creatinine  $\geq 1.5$  mg/dL
- Diabetes mellitus
- Known 3-vessel CAD
- Cerebrovascular disease
- Peripheral arterial disease
- Current smoker
- Known LVEF  $< 50\%$
- Chronic systolic or diastolic heart failure
- SBP  $\geq 140$  (within the past 12 months)
- LDL  $\geq 130$  (within the past 12 months)

## NOT TAKING ANY OF THE FOLLOWING:

- Brilinta (Ticagrelor)
- Coumadin (Warfarin)
- Eliquis (Apixaban)
- Pradaxa (Dabigatran)
- Savaysa (Edoxaban)
- Xarelto (Rivaroxaban)

## FOR MORE DETAILS CONTACT:

Darren DeWalt, MD, MPH, Principal Investigator  
Chief, Division of General Medicine and Clinical Epidemiology  
919-445-6806 | darren\_dewalt@med.unc.edu

Kristie Thompson  
Project Manager | UNC Sheps Center  
kristie\_Thompson@unc.edu

